



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ing from behind, whereas the flight from the nest was toward the clearer space in front. The sitting posture was not one of absolute rest at any time, as the head was constantly in motion, so that no approach could be made without her knowledge. The flight from the nest seemed to be directly out of it, without any preliminaries. The weather was warm, yet she would remain on the nest from fifteen to twenty minutes, and in no instance was away more than two minutes, while I had her under observation. The male frequently appeared in the vicinity, but neither offered food or even deigned to alight on the same tree, yet birds which had a good claim in the neighborhood dared not approach very close, as the combined attack of these active birds always proved so distasteful that they invariably beat a hasty retreat.—EDWIN H. EAMES, *Bridgeport, Conn.*

Snake Skins in the Nests of *Myiarchus crinitus*.—The habit of the Great Crested Flycatcher of putting scraps of discarded snake skin in its nest is—so far as the nests which I have found—invariable. Nevertheless, in one instance, at Tamworth, New Hampshire, I found a nest with one egg in it but with no snake skin visible. I found it about 7 A.M. one beautiful day in early July, 1888. I touched the egg and handled the nest slightly. Shortly before sunset I looked a second time into the hollow limb where the nest was placed, and was much surprised, in fact somewhat startled, by what I saw. Forming a complete circle about the egg, resting, in fact, like a wreath upon the circumference of the nest cavity, was a piece of snake skin six or seven inches long. The part which had encased the head of the snake was at the front of the nest and was slightly raised. It may not be wise to found a theory upon a single fact, but from the moment I saw that newly acquired snake skin, placed as it was, I made up my mind that the Great Crested Flycatcher uses the skin to scare away intruders. When the full set of eggs was laid, I took them and the nest. Only remnants of snake skin remained in the rubbish of the nest. The large skin had been removed or torn to bits.

The following year the same hollow was again occupied by Great Crested Flycatchers. I found the nest on July 7. It contained four eggs, and some scraps of snake skin were in sight. All the eggs hatched on the morning of the 12th. On the intermediate days my visits to the nest were regular. During those days a number of larger scraps of skin were placed on the outer edge of the nest. Their position was changed almost daily. Once some were set up like a fence, and so was a hen's feather. The birds knew of my visits, and scolded me while I remained in sight. These facts tended to confirm my theory in my own mind.—FRANK BOLLES, *Cambridge, Mass.*

Wintering of the Red-winged Blackbird near Cambridge, Mass.—On the 29th of December, 1889, while passing along the edge of a small swamp grown up with cat-tails, low bushes, birches, and maples, not far from Fresh Pond, I heard repeatedly the note of a Blackbird. I was un-